

Pain relief medication

Pain relief medication can help you to reduce the pain allowing you to undertake any suggested exercises and movements of the injury. Moving will help ease the pain and speed up your recovery significantly. Simple pain relief medications such as Paracetamol and Ibuprofen are often all that is required. Please read the medication instructions before taking anything.

Call 111 if you urgently need medical help or advice but it's not a life-threatening situation.

For less urgent health needs, contact your GP or local pharmacist in the usual way.

Minor Injury Unit

Sirona care & health's Minor Injury Unit offers treatment for adults, children and young people for a wide range of minor injuries. You can drop in without an appointment, or you may be referred by your GP or other healthcare professional. We aim to treat you within two hours.

Yate Minor Injuries Unit

Opening Times:
Open 7 days a week.
08.00 – 20.00

Address:
Yate West Gate Centre
21 West Walk
Yate Town Shopping Centre
BS37 4AX



For more information / Accessibility

t: 0300 124 5300*

This leaflet can be provided in other formats and languages, please contact us for more information.

e: askSirona@sirona-cic.org.uk

w: www.sirona-cic.org.uk



*Calls from landlines are charged up to 10p per minute; calls from mobiles vary, please check with your network provider. This is not a premium-rate number.

Date of creation: April 2019. Date of review: April 2021.
URN: 114

Registered Office:
Sirona care & health CIC, 2nd Floor, Kingswood Civic Centre
High Street, Kingswood, Bristol BS15 9TR
Company Number: 07585003



How to Treat Your Shoulder Injury

Information for you
if you have attended
the Minor Injury Unit

Service provided by

How to treat your injured shoulder

Following injury, your shoulder may be swollen, bruised and painful due to sprained muscles, tendons and ligaments. In order to help the natural healing process, follow the advice below.


During the first 3 days

- It is important to rest your arm to prevent further swelling. You may have been provided with a sling to support the arm. If so, use it.
- Frozen peas or crushed ice in a damp tea towel can be applied to the painful area. For maximum effect, apply for up to 20 minutes, every 2 hours. Cold can burn, so remove if uncomfortable.
- Once the worst of the pain begins to settle, it is important to start gently moving the injured arm. This may well be uncomfortable at first, but is essential to avoid future stiffness. Check that you can move your elbow, wrist and hand fully. Use your other arm to assist the movements of your injured shoulder. Move into discomfort, but short of pain.

Repeat 4 - 5 times daily.

Move your shoulder...

...forwards ...out to the side



After 3 days

You can now gradually discard your sling and progress your exercises to the following:

Exercise 1



Lean on a firm support with your uninjured arm. Allow your injured arm to hang loosely away from your body.

- a) Slowly swing your arm backwards and forwards alongside your body. Start with small movements and then gradually move as far as possible in each direction.
Repeat for 1-2 minutes.
- b) Now swing across your body and out to the side as far as possible.
Repeat for 1-2 minutes.
- c) Move the arm in a circular motion. Start with small circles increasing to larger circles.
Repeat for 1-2 minutes.

Exercise 2

Stand facing a wall, placing your hand flat against it. Slowly slide or 'walk' your hand up the wall as far as possible. You can use a piece of 'blu tack' to mark your daily progress. Remember to come down the wall slowly!

Repeat 2-3 times.

Exercise 2

Reach up behind your back as far as possible with your injured arm. Compare with your other arm.

Repeat 2-3 times.

Repeat exercises 4-5 times daily and, as your arm becomes more mobile, steadily build up your level of activity.

Depending on the nature and severity of your injury, it may take between 1-3 months, sometimes longer to recover. It is advisable to continue with the exercises until full movement has returned, although in some cases some permanent loss of movement may occur.

Problems that may persist for some time include pain at night, especially if lying on the injured shoulder. Also pain/discomfort when the shoulder is put in certain positions i.e. reaching behind your back or fully over your head.

Returning to sport

Before returning to racquet or contact sport, your shoulder should be free of pain, supple and strong. Consider weight training to build up your shoulder fitness. Just be sensible and steadily build up your exercise tolerance..